



CERTIFICATE OF ANALYSIS




Product Details

Product name:	Anti-4-1BB & FAP Reference Antibody (RO7122290)	Lot.No.:	P268000C
Target:	TNFRSF9 / 4-1BB / CD137, FAP	Catalog:	CHBA033
Target Accession:	Q07011 & P04626	Concentration :	1 mg/mL
Clonality:	Bispecific	Isotype:	IgG-VHH
Reactivity:	Human	Molecular Weight (kDa) :	177.78 kDa
Application:	Kinetics (SPR), ELISA, Bioactivity: FACS, Functional assay, Research in vivo	Endotoxin:	<1 EU/mg
Formulation:	Liquid: 100mM Pro-Ac, 20mM Arg pH 5.0 Lyophilization: 25mM histidine, 8% sucrose, 0.01% Tween80 pH6.2	Conjugation:	Unconjugated
Reconstitution:	For Powder, reconstitute with sterile, distilled water to a final concentration of 1 mg/ml. Gently shake to solubilize completely. Do not vortex.	Expression System:	CHO
Storage:	Upon receipt, store immediately at -20°C or lower for 24 months. Store aliquots at -80°C for up to 3 months. Avoid repeated freeze-thaw cycles.	Purification:	Protein A


Data

Purity: SDS-PAGE




Anti-4-1BB & HER2/neu Reference Antibody (Yh32367) on SDS-PAGE under reducing (R) condition. The purity of the protein is greater than 95%.

Purity: SEC-HPLC




The purity of Anti-4-1BB & HER2/neu Reference Antibody (Yh32367) is 98.99%, determined by SEC-HPLC.

ELISA



Yh32367 bound to 4-1BB protein, and then rebounded to secondary antibodies(Anti-Human-IgG-Fc-HRP), and read OD450. As shown in fig, Yh32367 bound to hu-4-1BB-His, and the EC50 was 0.007 nM.


ELISA



Yh32367 bound to Her2 protein, and then rebounded to secondary antibodies(Anti-Human-IgG-Fc-HRP), and read OD450. As shown in fig, Yh32367 bound to hu-Her2-His, and the EC50 was 0.009 nM.


The products are for research use only. Not for use in diagnostic procedures.

Bioactivity: FACS




Yh32367 bound to BT474 cells, and then rebounded to fluorescent secondary antibodies(Anti-Human IgG, Fcy PE) , and test by flow cytometry. As shown in fig, Yh32367 bound to BT474 cells, and the EC50 was 1.889 nM.

Bioactivity: FACS



Yh32367 bound to hu4-1BB-CHO-K cells, and then rebounded to fluorescent secondary antibodies(Anti-Human IgG, Fcy PE) , and test by flow cytometry. As shown in fig, Yh32367 bound to hu4-1BB-CHO-K cells, and the EC50 was 5.701 nM.

Function: Luciferase



Co-incubation of Yh32367 with BT474 cell and PBMCs for 4 hours, then LDH was detected to evaluate the ADCC activity of Yh32367. As shown in fig, Yh32367 has ADCC activity, and the IC50 was 0.054 nM.

The products are for research use only. Not for use in diagnostic procedures.